JOURNAL of ELECTROCHEMICAL SCIENCE and TECHNOLOGY

SUPPORTING INFORMATION

Binary and Ternary Solvents Influenced Growth of Thiol Staples Modified Nickel Nanoclusters and Their Tunable Supercapacitance

Ramadurai Murugan[1] | Vinitha Packirisamy[1] | Prabhu Pandurangan[2]*

[1]Center for Global Health Research, Saveetha Medical College and Hospitals, Saveetha Institute of Medical and Technical Science, Thandalam, Chennai- 602 105, Tamilnadu, India

[2]Department of Physical Chemistry, School of Chemical Science, University of Madras, Guindy Campus, Chennai, Tamilnadu-600 025, India



Fig. S1. Shows the UV-Visible absorption spectrum of [A] different ratios of methanol: water and [B] acetone: methanol: water isolated NiNCs respectively.



Fig. S2. Shows [A] the images of NiNCs different ratios of acetone, methanol, and water, [B] UV-Visible absorption spectrum of different ratio solvents and their soluble isolated NiO@Ni(II)SR-NCs respectively.

CORRESPONDENCE
T: +91 8939891487
E: pprabhumu@gmail.com
https://doi.org/10.33961/jecst.2025.01032

EPUB AHEAD OF PRINT

OPEN ACCESS This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/4.0) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.